

APPENDIX:

The Appendix includes the following items:

- ☒ - a Declaration
- ☒ - Exhibits 1-6

PATENT
0546-1082

IN THE U.S. PATENT AND TRADEMARK OFFICE

In re application of

Steve HEALD et al.

Conf. 4057

Application No. 10/550,945

Group 1651

Filed November 30, 2006

Examiner Irene Marx

**PREPARATION OF VANILLIN FROM MICROBIAL TRANSFORMATION MEDIA BY EXTRACTION BY
MEANS SUPERCRITICAL FLUIDS OR GASES**

DECLARATION

Assistant Commissioner for Patents

P.O. Box 1450

Alexandria, VA 22313-1450

Sir:

1. I, **Mylène DARRICAU**, am a French citizen. I am employed by the French firm SAFISIS (a subsidiary of LESAFFRE), in the capacity of R&D Project Manager, and have held that position since January 5th, 2009. My background is a Master of Engineering in Biotechnology with 7 years of experience in Fermentation and Microbiology; I have a diploma from Engineering School of INSA in Toulouse (France), and I am in charge of Research and Development projects for SAFISIS in Biomolecules.

I, **Evelyne FONCHY-PENOT**, hereby declare that I am a French citizen. I am employed by the French firm LESAFFRE INTERNATIONAL (a subsidiary of LESAFFRE), in the capacity of R&D Project Manager, and have held that position since 2000. LESAFFRE INTERNATIONAL is a service company which supports the different subsidiaries of LESAFFRE particularly in Research projects.

My background is PhD in biotechnology with more than 10 years experience in fermentation and Microbiology; I have a diploma from Engineer School from ESIL in Marseille (France), and I am in charge of Research and Development projects for LESAFFRE INTERNATIONAL in Biomolecules and Bioprocess.

2. We are familiar with the above-identified U.S. patent application, its prosecution before the United States Patent and Trademark Office, and the applied references of Rabenhorst et al. (U.S. Patent No. 6,133,003), Muheim et al. (U.S. Patent No. 6,235,507) and Makin (U.S. Patent No. 4,474,994).

3. In order to demonstrate the patentability of the present invention, we are submitting the following observations.

We confirm hereby that we have conducted the following experiments.

We state that, since my company got reports from independent laboratories that compare DSM 9991 and DSM 9992 of H&R patent and conclude that these two strains cannot be distinguished from one another; experiments were conducted on the sole DSM 9992 strain, exactly as disclosed in the H&R patent Examples 1 and 3.

We, Mrs DARRICAU and Mrs FONCHY-PENOT, who are researchers at SAF-ISIS and LESAFFRE INTERNATIONAL respectively, have realized the works detailed in the enclosed report (Exhibit 1)

As a summary the conclusions are that:

Strains DSM 9992 from US Patent No. 6,133,003 and Zyl 926 from the SAF-ISIS present application do not react at all the same way when applying, the same process as disclosed in examples 1 and 3 of US Patent No. 6,133,003.

Moreover, the HPLC analysis controls show that the ferulic acid was accumulated in the wort so that no bioconversion occurred with any of these two strains.

This means that the process in the US Patent No. 6,133,003 is inoperatively disclosed in order to get the alleged results regarding vanillin production.

As a consequence we state that the prior art was duly worked out with the intention to reduce to practice the working examples of US Patent No. 6,133,003, with the above mentioned absence of results.

We state that we made all reasonable due efforts, that a person of ordinary skill in the art should make, by taking all the information given in the specification such as for instance pH and oxygen maintenance, and also by using our general knowledge for example, to control the foaming that occurred during the fermentation by adding an antifoam.

Accordingly, we consider that, contrary to the DSM 9991 and 9992 strains of US Patent No. 6,133,003, the Zyl 926 microorganism, of the present application of SAF-ISIS, deposited within CABI as IMI 390106 for the purposes of patent procedures under the Budapest Treaty, provides unexpected results.

The latter are demonstrated notably by:

- a consumption of ferulic acid with concomitant production of above 11 g/L of vanillin
- a vanillin product obtained there from absent of odoriferous by-products and with none of guaiacol, vinyl guaiacol, eugenol and isoeugenol being present at more than 100ppm.

4. We declare that all statements made herein of our own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under §1001 of Title 18 of the United States code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Date June 29, 2010

Myliène DARRICAU
R&D Product Manager
SAF-ISIS
Zone Artisanale
Soustons 40140 – France



Evelyne FONCHY-PENOT
R&D Project Manager
LESAFFRE INTERNATIONAL
Rue Gabriel Peri 147
Marcq en Baroeul 59700 – France




Exhibit 1

Report on the “Characterisation of Strains belonging to the Genus *Amycolatopsis* by 16S rRNA Gene Sequencing”

Professor M. Goodfellow

Background

Initially, the client wished to have the taxonomic relationships between the three *Amycolatopsis* strains determined by 16S rRNA gene sequencing. This study was designed to establish the inferred phylogenetic relationships between the cultures and their position (s) in the *Amycolatopsis* 16S rRNA gene tree.

Receipt of Strains

The package containing the strains arrived intact on May 20th (See photographs attached).

Materials and Methods

Cultivation of strains. The strains were grown in shake flasks (150 rpm) in ISP 2 broth for 48 hours at 37°C. Following incubation, biomass was harvested by centrifugation.

16S rRNA gene sequencing analyses. Chromosomal DNA was isolated and 16S rRNA PCR was done using well established procedure (Lane, 1991). PCR product was purified and direct sequencing of 16S rRNA genes of the three *Amycolatopsis* strains was carried out using BigDye™ version 3.1 Terminator Cycle Sequencing Kit (PE Applied Biosystems, MA). Samples were completed using 10 ng of DNA for every 100 bp of template DNA and 3.2 pmol of primer. Post-sequencing reactions were purified away from unincorporated dye terminators using Performa DTR spin-columns (EdgeBiosystems, MD). Sequences were determined using an ABI 3730 Sequencer. The resultant almost complete 16S rRNA gene sequences (1360 nucleotides) were aligned manually against corresponding sequences of the type strains of *Amycolatopsis* species, retrieved from GenBank, using PHYDIT software (Chun, 1995). A phylogenetic tree was inferred by using neighbour-joining (Saitou & Nei, 1987) tree-making algorithm from the PHYDIT program and an evolutionary distance matrix generated by using the distance model of Jukes & Cantor (1969). The robustness of the phyletic lines recovered in the *Amycolatopsis* tree were determined in a bootstrap analysis (Felsenstein, 1985) of the neighbour-joining dataset by using the SEQBOOT and CONSENSE options from the PHYLIP suite of programs (Felsenstein, 1993).

Results

It is apparent from Figures 1 to 3 that the 16S rRNA gene sequences obtained for the three *Amycolatopsis* strains are of high quality as there are no ambiguities in the conserved regions of the traces. It is clear from the 16S rRNA *Amycolatopsis* tree that the three strains belong to the *Amycolatopsis methanolica* subclade, the taxonomic integrity of the latter is underpinned by a 100% bootstrap value (Fig. 4). It is also apparent from the tree and from the associated nucleotide similarity and difference

matrix (Table 1) that the three strains share identical 16S rRNA gene sequences with one another.

Conclusions

- The three *Amycolatopsis* strains belong to the *Amycolatopsis methanolica* subclade in the 16S rRNA *Amycolatopsis* gene tree.
- The three *Amycolatopsis* strains shared identical 16S rRNA gene sequences with one another.

References

Chun, J. (1995). Computer-assisted classification and identification of actinomycetes. Ph.D. Thesis, University of Newcastle, Newcastle upon Tyne, UK.

Felsenstein, J. (1985). Confidence limits on phylogeny: An appropriate use of the bootstrap. *Evolution* **39**, 7783-7791.

Felsenstein, J. (1993). PHYLIP- Phylogenetic Inference Package version 3.5.1. Department of Genetics, University of Washington, Seattle, USA.

Jukes, T.H. & Cantor, C.R. (1969). Evolution of protein molecules. In *Mammalian Protein Metabolism*, volume **3**, pp. 21-132. Edited by H. N. Munro, Academic Press, New York.

Lane, D.J. (1991). 16S/23S rRNA Sequencing. In *Nucleic Acid Techniques in Bacterial Systematics*, pp. 115-148. Edited by E. Stackebrandt & M. Goodfellow. John Wiley & Sons, Chichester, New York.

Saitou, N. & Nei, M. (1987). The neighbour-joining method: a new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution* **4**, 406-425.

Attachments with the Report

Legends for Figures.

Figs. 1a-3b. Chromatograms of Samples A, B and C respectively (a- sequencing with forward primer; b- sequencing with reverse primer).


Fig. 4. Neighbour-joining tree based on nearly complete 16S rRNA gene sequences showing relationships between the three *Amycolatopsis* strains and between them and the type strains of *Amycolatopsis* species. Numbers at the nodes indicate the level of bootstrap support (%) based on neighbour-joining analysis of 1000 resampled datasets; only values above 50% are given. The scale bar indicates 0.02 substitutions per nucleotide position.

Photographs.

Photographs 1-5. Receipt of strains.

Tables.

Table 1. Nucleotide similarity and difference matrix with the three strains which share identical 16S rRNA gene sequences with one another.

A handwritten signature in black ink, appearing to read "M. Goodfellow". The signature is written in a cursive, flowing style.

Professor Michael Goodfellow

CCFRA Technology Limited
Station Road, Chipping Campden
Gloucestershire
GL55 6LD, United Kingdom

Tel: +44 (0)1386 842000
Fax: +44 (0)1386 842100
www.campden.co.uk

Richard Tailon
LeSaffre Saffris
Zone Artisanale
40140 Soustons
France

Exhibit 2



Campden BRI 
food and drink innovation

Report No: MB/REP/MMR/115732/00002-00004/2
Report Date: 30/06/09

CONFIDENTIAL
CERTIFICATE OF ANALYSIS
Microbiology Department
MicroID

Notes

The results provided herein relate only to the items tested.

RiboPrint[®], RiboGroup[®] and RiboPrinter[™] are trademarks of Qualicon[™] L.L.C., a subsidiary of E.I. DuPont de Nemours and Company, Wilmington, Delaware, USA.

Prepared/Checked By:

Authorised By:

Authorised	C. Mitchell	
	Research Technical Officer	
Signatories:	S. Jordan	
	Molecular Methods Manager	
	C. Baylis	/
	Methods Research Manager	

Any opinions and interpretations contained in this report are outside the scope of UKAS accreditation.

This test report or certificate shall not be reproduced, except in full, without written approval of Campden BRI.

Confidentiality notice: The information in this document is only intended for the individual or entity to whom it is addressed. It may contain privileged and confidential information that is exempt from disclosure by law and if you are not the intended recipient, you must not copy, distribute or take any action in reliance on it. If you have received this document in error, please notify Campden BRI immediately by telephone.

Report No: MB/REP/MMR/115732/00002-00004/2

Page 1 of 6

CCFRA Technology Limited, Registered No. 3836922, Incorporated in
England & Wales
Registered Office: Station Road, Chipping Campden, Glos. GL55 6LD
Part of Campden BRI Group

Information emanating from Campden BRI is given after the exercise of all
reasonable care and skill in its compilation, preparation and issue, but is
provided without liability in its application and use.

Summary

Three bacterial isolates were sent to Campden BRI from SAFISIS for strain comparison using the Qualicon™ RiboPrinter® at the request of the client. The parcel was received on site unopened as shown in the images taken of the package in Appendix 1.

The isolates were stored and grown prior to analysis as detailed in the NDA supplied by the client. Standard techniques and reagents were used for the ribotyping. The enzyme PvuII was used for the analysis of the isolates as it has been shown to differentiate between DuPont database entries for *Streptomyces* sp. The results of the work are noted in Tables below, with a pattern comparison shown in Figure 1. A summary of the RiboPrinter methodology is outlined in Appendix 2.

CCFRA Code:	MMR/115732/00002	Date of Receipt:	20/05/09
Customer Code	Sample A	Client presumptive identification	<i>Amycolatopsis</i> sp
Sample Description:	Frozen culture		
Comments:	Stored at ambient temperature prior to analysis		
Analysis Type	Test Details	Identification	Comments
RiboPrinter®	PvuII	None	Isolate fell outside of 0.85 similarity needed for automatic identification against the DuPont identification database. The isolate was placed in RiboGroup 114-3210-S-1.
RiboPrinter®	Strain comparison		Refer to Figure 1 for details

CCFRA Code:	MMR/115732/00003	Date of Receipt:	20/05/09
Customer Code	Sample B	Client presumptive identification	<i>Amycolatopsis</i> sp
Sample Description:	Freeze Dried Ampoule		
Comments:	Stored at ambient temperature prior to analysis		
Analysis Type	Test Details	Identification	Comments
RiboPrinter®	PvuII	None	Isolate fell outside of 0.85 similarity needed for automatic identification against the DuPont identification database. The isolate was placed in RiboGroup 114-3210-S-1.
RiboPrinter®	Strain comparison		Refer to Figure 1 for details

CCFRA Code:	MMR/115732/00004	Date of Receipt:	20/05/09
Customer	Sample C	Client presumptive	<i>Amicolatopsis</i> sp
Code		Identification	
Sample	Freeze Dried Ampoule		
Description:			
Comments:	Stored at ambient temperature prior to analysis		
Analysis Type	Test Details	Identification	Comments
RiboPrinter®	<i>PvuII</i>	None	Isolate fell outside of 0.85 similarity needed for automatic identification against the DuPont identification database. The isolate was placed in RiboGroup 114-3210-S-1.
RiboPrinter®	Strain comparison		Refer to Figure 1 for details

Strain comparison

A comparison of the patterns generated for the three isolates is shown below in Figure 1a and b.

Figure 1a: Strain comparison of isolates MB/115732/2-4 using MB/115732/2 as the reference isolate





Figure 1b: Strain comparison of isolates MB/115732/2-4 using MB/115732/3 as the reference isolate




All three isolates were placed in the same Ribogroup 114-3210-S-1. These results suggest that the isolates are indistinguishable from each other using PvuII.

These data readily demonstrate the ability of the RiboPrinter system to characterise and discriminate bacterial strains by analysis of RiboPrint patterns. This information will be available for reference on the archived database and as such can be used to authenticate, or compare to, these cultures time and time again.

These isolates will be kept for a standard period of 4 weeks after ribotyping after which they will be destroyed as outlined in the NDA supplied by the client.

Appendix 1: Image of package received for analysis at Campden BRI



The RiboPrinter system is an automated ribotyping process as follows:

Bacterial cells are broken open to release the genomic DNA into solution. This DNA is digested ('cut up') using a restriction enzyme to generate fragments of different lengths, which are then separated out on the basis of size. Ribosomal DNA is detected using probes which give off light that is captured by a camera. The resulting image is converted into a RiboPrint pattern that can be characterised (into RiboGroups) and where possible identified by proprietary software. RiboGroups are assigned to patterns in the following format - RiboGroup library - Machine number - batch number and sample position where the pattern occurred for the first time for example RIBO1 114-1002-S-1

As each sample is analysed by the software, it makes a decision about how closely the new sample pattern matches the existing patterns in the database. If the software finds a match, that is if it is unable to distinguish the new pattern from an existing pattern (i.e. >92% similarity between patterns), it assigns the new sample to that RiboGroup. All samples in a RiboGroup have the same RiboGroup code. Thus if the isolate has been previously ribotyped on this instrument, the instrument database will recognise the reproduced RiboPrint and allocate the RiboGroup code assigned in the first instance.

When an isolate is analysed on the machine for the first time, i.e. it does not match with any existing patterns, it will receive a new code i.e. the batch and sample number specific to that new batch run.

For automatic (DuPont) identification, the new sample information is compared to existing patterns within a supplied (DUP) database. If the software finds a match of >85% a DuPont identification will be reported.

If no automatic identification is given, the user can instruct the software to show the percentage similarities between existing patterns in the instrument sample database (Campden BRI database) and the new sample pattern. When >85% similarity is shown identification can be offered.

Exhibit 3

PHYDIT - Similarity analysis
 1360 nucleotides analysed
 Lower-left triangle contains [NT] Similarity.
 Uppper-right triangle contains [NT] Different/Total nucleotides.

The following table contains tab-delimited numbers. Copy and paste to MS Excel or Word.

	Amycolatopsis Sample A
Amycolatopsis Sample A	---
Amycolatopsis Sample B	100
Amycolatopsis Sample C	100
Amycolatopsis methanolica DSM 44096T (AJ249135)	98.88
Amycolatopsis thermoflava DSM 44574T (AF052390)	98.88
Amycolatopsis jejuensis NRRL B-24427T (DQ000200)	94.25
Amycolatopsis sulphurea DSM 46092T (AJ293756)	95.37
Amycolatopsis alba DSM 44262T (AF051340)	94.76
Amycolatopsis coloradensis DSM 44225T (AJ293753)	94.47
Amycolatopsis regifaucium DSM 45072T (AY129760)	94.84
Amycolatopsis azurea DSM 43854T (AJ400709)	94.69
Amycolatopsis orientalis DSM 40040T (AJ400711)	94.69
Amycolatopsis keratiniphila subsp. keratiniphila DSM 4440T (AJ278496)	94.69
Amycolatopsis keratiniphila subsp. nogabecina DSM 445T (AJ508238)	94.69
Amycolatopsis japonica DSM 44213T (AJ508236)	94.54
Amycolatopsis decaplanina DSM 44594T (AJ508237)	94.77
Amycolatopsis minnesotensis NRRL B-24435 (DQ076482)	94.84
Amycolatopsis nigrescens DSM 44992T (DQ486888)	95.44
Amycolatopsis lexingtonensis DSM 44653T (AY183358)	94.61
Amycolatopsis pretoriensis DSM 44654T (AY183356)	94.54
Amycolatopsis rifamycinica DSM 46095T (AY083603)	94.69
Amycolatopsis kentuckyensis DSM 44652T (AY183357)	94.54
Amycolatopsis bathimycina DSM 44591T (AJ508239)	94.47
Amycolatopsis vancoresmycina DSM 44592T (AJ508240)	94.84
Amycolatopsis plumensis DSM 44776T (AY262825)	94.61
Amycolatopsis tolypomycina DSM 44544T (AJ293757)	94.61
Amycolatopsis mediterranei DSM 43304T (AJ293754)	94.84
Amycolatopsis australiensis DSM 44671T (AY129753)	95.14
Amycolatopsis saalfeldensis DSM 44993T (DQ792500)	95.07
Amycolatopsis echigonensis JCM 21831T (AB248535)	94.84
Amycolatopsis nilgatensis JCM 21832T (AB248537)	95.06
Amycolatopsis benzoatilytica DSM 43387T (AY957506)	93.39
Amycolatopsis albidoflavus DSM 44639T (AJ252832)	94.9
Amycolatopsis halotolerans NRRL B-24428T (DQ000196)	95.21
Amycolatopsis rubida DSM44637T (AF222022)	94.51
Amycolatopsis sacchari DSM 44468T (AF223354)	95.29
Amycolatopsis taiwanensis NBRC 102103T (DQ160215)	94.99
Amycolatopsis marina NBRC 104263T (EU329845)	95.96
Amycolatopsis palatopharyngis JCM 12460T (AF479268)	96.19
Amycolatopsis fastidiosa DSM 43855T (AB184566)	92.38

Amycolatopsis Sample B	Amycolatopsis Sample C
0/1338	0/1338
---	0/1338
100 ---	
98.88	98.88
98.88	98.88
94.25	94.25
95.37	95.37
94.76	94.76
94.47	94.47
94.84	94.84
94.69	94.69
94.69	94.69
94.69	94.69
94.69	94.69
94.54	94.54
94.77	94.77
94.84	94.84
95.44	95.44
94.61	94.61
94.54	94.54
94.69	94.69
94.54	94.54
94.47	94.47
94.84	94.84
94.61	94.61
94.61	94.61
94.84	94.84
95.14	95.14
95.07	95.07
94.84	94.84
95.06	95.06
93.39	93.39
94.9	94.9
95.21	95.21
94.51	94.51
95.29	95.29
94.99	94.99
95.96	95.96
96.19	96.19
92.38	92.38

Amycolatopsis methanolica DSM 44096T (AJ249135)

15/1338

15/1338

15/1338

99.85

94.32

94.99

94.69

94.47

94.39

94.62

94.54

94.77

94.61

94.47

94.92

94.54

95.74

94.61

94.54

94.62

94.47

94.39

94.47

94.17

94.24

94.76

94.99

95.22

94.54

94.76

92.93

94.75

94.91

94.81

95.74

95.44

96.34

96.49

92.38

Amycolatopsis thermoflava DSM 44574T (AF052390)

15/1337

15/1337

15/1337

2/1337

94.24

94.99

94.61

94.39

94.31

94.54

94.47

94.69

94.54

94.39

94.84

94.47

95.66

94.54

94.47

94.54

94.39

94.32

94.39

94.09

94.16

94.68

94.91

95.14

94.46

94.68

92.85

94.67

94.83

94.73

95.66

95.36

96.26

96.41

92.3

Amycolatopsis jejuensis NRRL B-24427T (DQ000200)

77/1338

77/1338

77/1338

76/1338

77/1337

97.83

96.86

96.64

97.08

97.01

96.79

96.64

96.86

97.01

96.79

96.71

96.71

96.56

96.64

96.64

96.41

96.19

96.64

96.34

96.78

96.48

97.01

97.16

97.23

97.16

95.29

97

96.86

96.77

96.64

93.88

95.44

95.52

90.44

Amycolatopsis sulphurea DSM 46092T (AJ293756)	Amycolatopsis alba DSM 44262T (AF051340)
62/1338	70/1337
62/1338	70/1337
62/1338	70/1337
67/1338	71/1337
67/1337	72/1336
29/1339	42/1337
---	45/1337
	96.63 ---
	96.49
	97.38
	97.01
	96.94
	96.79
	97.01
	96.94
	97.31
	97.01
	97.09
	96.63
	96.56
	96.56
	96.34
	96.26
	96.71
	96.11
	96.41
	96.41
	97.09
	97.16
	97.31
	97.68
	95.67
	97.68
	97.68
	96.92
	97.23
	93.58
	95.52
	95.67
	91.04
	99.25
	98.28
	98.65
	98.58
	98.43
	98.5
	98.28
	98.35
	97.38
	97.53
	97.83
	97.91
	97.98
	97.75
	97.68
	97.68
	97.16
	97.6
	97.9
	97.46
	97.83
	97.38
	97.53
	95.82
	97.6
	97.53
	97.22
	96.19
	94.09
	95.14
	95.44
	91.1

Amycolatopsis coloradensis DSM 44225T (AJ293753)

74/1338

74/1338

74/1338

74/1338

75/1337

45/1338

47/1338

10/1337

98.5

98.58

98.8

98.65

98.65

98.65

98.58

97.61

97.38

97.83

97.91

97.83

97.61

97.68

97.46

97.08

97.53

97.68

97.46

97.91

97.08

97.23

95.59

97.45

97.23

97.14

95.89

93.87

95.07

95.37

90.81

Amycolatopsis regifaucium DSM 45072T (AY129760)

69/1337

69/1337

69/1337

75/1337

76/1336

39/1337

35/1337

23/1336

20/1337

98.88

99.1

98.88

99.25

99.48

99.1

97.46

97.01

97.46

97.38

97.38

97.16

97.01

97.46

97.01

97.38

97.3

97.83

97.23

97.83

98.13

96.43

98.05

98.13

97.29

95.96

93.34

95.29

95.29

91.17

Amycolatopsis azurea DSM 43854T (AJ400709)	Amycolatopsis orientalis DSM 40040T (AJ400711)
71/1338	71/1338
71/1338	71/1338
71/1338	71/1338
72/1338	73/1338
73/1337	74/1337
40/1338	43/1338
40/1338	41/1338
18/1337	19/1337
19/1338	16/1338
15/1337	12/1337
---	6/1338
	99.55 ---
	99.18
	99.48
	99.1
	99.18
	99.33
	97.16
	97.31
	97.31
	98.13
	97.83
	98.06
	97.76
	97.91
	97.61
	97.68
	97.38
	97.68
	97.38
	97.53
	97.38
	96.93
	96.86
	97.46
	97.38
	97.75
	97.46
	97.76
	97.76
	97.46
	97.61
	97.38
	97.9
	97.68
	96.43
	96.35
	97.9
	97.83
	97.75
	97.68
	97.67
	97.44
	96.26
	96.26
	93.87
	93.65
	95.74
	95.59
	96.04
	95.81
	91.63
	91.63

Amycolatopsis keratiniphila subsp. keratiniphila DSM 4440T (AJ278496)

71/1338
71/1338
71/1338
70/1338
71/1337
45/1338
43/1338
21/1337
18/1338
15/1337
11/1338
9/1338

99.7
99.25
99.33
97.31
97.53
97.83
97.76
97.46
97.23
97.23
97.46
96.86
97.31
97.38
97.61
97.38
97.76
97.9
96.05
97.98
97.75
97.89
96.49
93.87
95.89
96.11
91.55

Amycolatopsis keratiniphila subsp. nogabecina DSM 445T (AJ508238)

71/1337
71/1337
71/1337
72/1337
73/1336
42/1337
40/1337
20/1336
18/1337
10/1336
7/1337
5/1337
4/1337

99.33
99.48
97.31
97.38
97.75
97.68
97.53
97.31
97.31
97.46
96.78
97.31
97.38
97.76
97.38
97.83
98.13
96.43
98.05
97.98
97.89
96.41
93.79
95.74
95.96
91.77

Amycolatopsis japonica DSM 44213T (AJ508236)	Amycolatopsis decaplanina DSM 44594T (AJ508237)	
73/1338	70/1338	
73/1338	70/1338	
73/1338	70/1338	
74/1338	68/1338	
75/1337	69/1337	
40/1338	43/1338	
41/1338	36/1338	
23/1337	22/1337	
18/1338	19/1338	
7/1337	12/1337	
12/1338	11/1338	
11/1338	9/1338	
10/1338	9/1338	
9/1337	7/1337	
---	11/1338	
	99.18 ---	
	97.23	97.09
	97.16	97.23
	97.83	97.38
	97.76	97.31
	97.61	97.31
	97.38	97.08
	97.23	96.94
	97.31	97.09
	96.93	96.48
	97.46	96.93
	97.46	97.31
	97.68	97.98
	97.38	97.09
	97.53	97.68
	97.75	98.13
	95.9	96.12
	97.75	97.98
	97.6	97.98
	97.37	97.67
	95.96	96.19
	93.57	93.57
	95.52	95.89
	95.59	95.81
	91.11	91.41

Amycolatopsis minnesotensis NRRL B-24435 (DQ076482)

69/1338

69/1338

69/1338

73/1338

74/1337

44/1339

40/1339

35/1337

32/1338

34/1337

38/1338

36/1338

36/1338

36/1337

37/1338

39/1338

98.43

96.71

96.79

96.64

96.41

96.56

97.38

96.26

96.56

96.63

96.94

97.76

96.41

96.56

95.14

96.78

96.56

96.24

96.86

93.13

95.96

96.11

91.04

Amycolatopsis nigrescens DSM 44992T (DQ486888)

61/1338

61/1338

61/1338

57/1338

58/1337

44/1339

39/1339

33/1337

35/1338

40/1337

36/1338

36/1338

33/1338

35/1337

38/1338

37/1338

21/1339

96.71

96.79

96.79

96.56

96.49

96.64

96.04

96.41

96.78

97.09

97.98

96.49

96.64

94.84

96.78

96.64

96.7

97.53

93.5

96.64

96.64

91.78

Amycolatopsis lexiingtonensis DSM 44653T (AY183358)

72/1337

72/1337

72/1337

72/1337

73/1336

46/1337

45/1337

29/1336

29/1337

34/1336

25/1337

29/1337

29/1337

30/1336

29/1337

35/1337

44/1337

44/1337

99.93

99.63

99.4

99.25

98.95

98.8

99.1

99.1

98.28

98.5

97.6

97.75

96.65

97.97

97.75

97.59

96.34

94.32

95.36

95.51

91.47

Amycolatopsis pretoriensis DSM 44654T (AY183356)

73/1338

73/1338

73/1338

73/1338

74/1337

45/1339

46/1339

28/1337

28/1338

35/1337

26/1338

30/1338

30/1338

31/1337

30/1338

36/1338

43/1339

43/1339

1/1337

99.7

99.48

99.33

99.03

98.88

99.18

99.18

98.36

98.58

97.53

97.68

96.58

97.9

97.68

97.52

96.41

94.4

95.29

95.44

91.41

Amycolatopsis rifamycinica DSM 46095T (AY083603)

71/1338

71/1338

71/1338

72/1338

73/1337

45/1338

46/1338

27/1337

29/1338

35/1337

28/1338

32/1338

34/1338

33/1337

32/1338

36/1338

45/1338

43/1338

5/1337

4/1338

99.63

99.18

99.03

98.65

98.95

99.33

98.36

98.43

97.68

97.83

96.43

97.9

97.83

97.52

96.34

94.32

95.37

95.37

91.41

Amycolatopsis kentuckyensis DSM 44652T (AY183357)

73/1337

73/1337

73/1337

74/1337

75/1336

48/1337

49/1337

30/1336

32/1337

38/1336

31/1337

35/1337

37/1337

36/1336

35/1337

39/1337

48/1337

46/1337

8/1336

7/1337

5/1337

99.1

98.95

98.58

98.88

99.25

98.28

98.2

97.46

97.6

96.2

97.67

97.75

97.44

96.04

94.02

95.21

95.21

91.1

Amycolatopsis balhimycina DSM 44591T (AJ508239)

74/1338

74/1338

74/1338

75/1338

76/1337

51/1338

50/1338

31/1337

31/1338

40/1337

31/1338

35/1338

37/1338

36/1337

37/1338

41/1338

46/1338

47/1338

10/1337

9/1338

11/1338

12/1337

98.65

98.35

98.65

99.18

98.28

98.36

97.16

97.31

96.2

97.53

97.46

97.29

95.96

94.17

94.92

95.07

91.11

Amycolatopsis vancoresmycina DSM 44592T (AJ508240)

69/1338

69/1338

69/1338

74/1338

75/1337

45/1338

44/1338

31/1337

34/1338

34/1337

33/1338

35/1338

34/1338

34/1337

36/1338

39/1338

35/1338

45/1338

14/1337

13/1338

13/1338

14/1337

18/1338

98.65

98.95

98.65

98.36

98.13

97.91

98.05

96.66

98.13

98.2

97.59

96.26

94.17

95.29

95.29

91.33

Amycolatopsis plumensis DSM 44776T (AY262825)

72/1337

72/1337

72/1337

78/1337

79/1336

49/1337

52/1337

38/1336

39/1337

40/1336

41/1337

42/1337

42/1337

43/1336

41/1337

47/1337

50/1337

53/1337

16/1336

15/1337

18/1337

19/1336

22/1337

18/1337

99.1

98.28

97.83

97.46

97.38

97.45

96.27

97.45

97.6

96.84

95.74

93.79

94.76

94.99

90.8

Amycolatopsis tolypomycina DSM 44544T (AJ293757)

72/1337

72/1337

72/1337

77/1337

78/1336

43/1337

48/1337

32/1336

33/1337

35/1336

34/1337

35/1337

36/1337

36/1336

34/1337

41/1337

46/1337

48/1337

12/1336

11/1337

14/1337

15/1336

18/1337

14/1337

12/1336

98.58

98.2

97.91

97.6

97.75

96.58

97.82

97.9

97.14

95.89

94.02

95.06

95.36

91.1

Amycolatopsis mediterranei DSM 43304T (AJ293754)

69/1336

69/1336

69/1336

70/1336

71/1335

47/1336

48/1336

28/1335

31/1336

36/1335

30/1336

34/1336

35/1336

35/1335

34/1336

36/1336

45/1336

43/1336

12/1335

11/1336

9/1336

10/1335

11/1336

18/1336

23/1335

19/1335

98.5

98.35

97.38

97.53

96.12

97.6

97.68

97.36

96.11

94.01

95.36

95.36

91.39

Amycolatopsis australiensis DSM 44671T (AY129753)

65/1338

65/1338

65/1338

67/1338

68/1337

40/1338

39/1338

34/1337

34/1338

29/1337

30/1338

30/1338

32/1338

30/1337

31/1338

27/1338

41/1338

39/1338

23/1337

22/1338

22/1338

23/1337

23/1338

22/1338

29/1337

24/1337

20/1336

97.91

97.46

97.75

96.35

97.98

97.9

97.52

96.71

94.25

95.96

95.81

91.41

Amycolatopsis saalfeldensis DSM 44993T (DQ792500)

66/1338

66/1338

66/1338

64/1338

65/1337

38/1338

38/1338

29/1337

28/1338

37/1337

30/1338

34/1338

35/1338

35/1337

35/1338

39/1338

30/1338

27/1338

20/1337

19/1338

21/1338

24/1337

22/1338

25/1338

34/1337

28/1337

22/1336

28/1338

97.16

97.31

95.74

97.53

97.31

97.22

97.01

94.39

95.74

95.89

91.48

Amycolatopsis echigonensis JCM 21831T (AB248535)

69/1337

69/1337

69/1337

73/1337

74/1336

37/1338

36/1338

35/1336

39/1337

29/1336

32/1337

35/1337

30/1337

29/1336

33/1337

31/1337

48/1338

47/1339

32/1336

33/1338

31/1337

34/1336

38/1337

28/1337

35/1336

32/1336

35/1335

34/1337

38/1337

99.4

97.19

98.95

99.1

98.5

96.19

94.02

95.44

95.29

90.88

Amycolatopsis niigatensis JCM 21832T (AB248537)

66/1336

66/1336

66/1336

70/1336

71/1335

38/1337

31/1337

33/1335

37/1336

25/1335

28/1336

31/1336

28/1336

25/1335

30/1336

25/1336

46/1337

45/1338

30/1335

31/1337

29/1336

32/1335

36/1336

26/1336

34/1335

30/1335

33/1334

30/1336

36/1336

8/1338

97.57

99.18

99.55

98.42

96.18

93.94

95.43

95.21

91.25

Amycolatopsis benzoatilytica DSM 43387T (AY957506)

87/1316

87/1316

87/1316

93/1316

94/1315

62/1316

57/1316

55/1315

58/1316

47/1315

47/1316

48/1316

52/1316

47/1315

54/1316

51/1316

64/1316

68/1317

44/1315

45/1316

47/1316

50/1315

50/1316

44/1316

49/1315

45/1315

51/1314

48/1316

56/1316

37/1316

32/1316

97.41

97.42

96.64

94.53

92.17

93.54

93.62

90.05

Amycolatopsis albidoflavus DSM 44639T (AJ252832)

68/1334

68/1334

68/1334

70/1334

71/1333

40/1335

31/1335

32/1333

34/1334

26/1333

28/1334

29/1334

27/1334

26/1333

30/1334

27/1334

43/1335

43/1336

27/1333

28/1335

28/1334

31/1333

33/1334

25/1334

34/1333

29/1333

32/1332

27/1334

33/1334

14/1336

11/1336

34/1314

—

99.18

98.5

96.33

93.93

95.35

95.28

91.32

Amycolatopsis halotolerans NRRL B-24428T (DQ000196)

64/1336

64/1336

64/1336

68/1336

69/1335

42/1337

31/1337

33/1335

37/1336

25/1335

30/1336

31/1336

30/1336

27/1335

32/1336

27/1336

46/1337

45/1338

30/1335

31/1337

29/1336

30/1335

34/1336

24/1336

32/1335

28/1335

31/1334

28/1336

36/1336

12/1338

6/1338

34/1316

11/1336

98.42

96.18

93.72

95.28

95.21

91.25

Amycolatopsis rubida DSM44637T (AF222022)	Amycolatopsis sacchari DSM 44468T (AF223354)
73/1330	63/1338
73/1330	63/1338
73/1330	63/1338
69/1330	57/1338
70/1329	58/1337
43/1331	45/1338
41/1331	37/1338
37/1329	51/1337
38/1330	55/1338
36/1329	54/1337
31/1330	50/1338
34/1330	50/1338
28/1330	47/1338
28/1329	48/1337
35/1330	54/1338
31/1330	51/1338
50/1331	42/1338
44/1332	33/1338
32/1329	49/1337
33/1331	48/1338
33/1330	49/1338
34/1330	53/1337
36/1330	54/1338
32/1330	50/1338
42/1329	57/1337
38/1329	55/1337
35/1328	52/1336
33/1330	44/1338
37/1330	40/1338
20/1332	51/1337
21/1332	51/1336
44/1310	72/1316
20/1330	49/1334
21/1332	51/1336
---	46/1330
	96.54 ---
	93.99
	95.79
	95.64
	90.98
	94.77
	96.26
	96.41
	91.85

Amycolatopsis taiwanensis NBRC 102103T (DQ160215)

67/1338

67/1338

67/1338

61/1338

62/1337

82/1339

86/1339

79/1337

82/1338

89/1337

82/1338

85/1338

82/1338

83/1337

86/1338

86/1338

92/1339

87/1339

76/1337

75/1339

76/1338

80/1337

78/1338

78/1338

83/1337

80/1337

80/1336

77/1338

75/1338

80/1338

81/1337

103/1316

81/1335

84/1337

80/1331

70/1338

94.1

94.02

89.92

Amycolatopsis marina NBRC 104263T (EU329845)

54/1338

54/1338

54/1338

49/1338

50/1337

61/1338

60/1338

65/1337

66/1338

63/1337

57/1338

59/1338

55/1338

57/1337

60/1338

55/1338

54/1338

45/1338

62/1337

63/1338

62/1338

64/1337

68/1338

63/1338

70/1337

66/1337

62/1336

54/1338

57/1338

61/1337

61/1336

85/1316

62/1334

63/1336

56/1330

50/1338

79/1338

99.4

91.55

Amycolatopsis palatopharyngis JCM 12460T (AF479268)

51/1338

51/1338

51/1338

47/1338

48/1337

60/1338

58/1338

61/1337

62/1338

63/1337

53/1338

56/1338

52/1338

54/1337

59/1338

56/1338

52/1338

45/1338

60/1337

61/1338

62/1338

64/1337

66/1338

63/1338

67/1337

62/1337

62/1336

56/1338

55/1338

63/1337

64/1336

84/1316

63/1334

64/1336

58/1330

48/1338

80/1338

8/1342

91.78

Amycolatopsis fastidiosa DSM 43855T (AB184566)

102/1338
102/1338
102/1338
102/1338
103/1337
128/1339
120/1339
119/1337
123/1338
118/1337
112/1338
112/1338
113/1338
110/1337
119/1338
115/1338
120/1339
110/1339
114/1337
115/1339
115/1338
119/1337
119/1338
116/1338
123/1337
119/1337
115/1336
115/1338
114/1338
122/1338
117/1337
131/1316
116/1336
117/1337
120/1331
109/1338
135/1339
113/1338
110/1338

Exhibit 4

>a1-27F

GCTGAAGCACCTTACGGGTGTGGATGAGTGGCGAACGGGTGAGTAACACGTGGGTAACCT
GCCCCTGTACTTTGGGATAAGCCCTGGAAACGGGGTCTAATACCGAATATGACTCGCTAAG
GCATCTTTGGTGGGTGGGAAAGTTTCGGCGGTACAGGATGGGCCCCGGGCTATCAGCTTGT
TGGTGGGGTAGTGGCCTACCAAGCGACGACGGGTAGCCGGCCTGAGAGGGTGACCGGCC
ACACTGGGACTGAGACACGGGCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGCAC
AATGGGCGGAAGCCTGATGCAGCGACGCCGCTGAGGGATGACGGCCTTCGGGTTGTAAA
CCTCTTTCCGCCAGGGACGAAGCGCAAGTGACGGTACCTGGAGAAGAAGCACCGGCTAACT
ACGTGCCAGCAGCCGCGGTAATACGTAGGGTGCAAGCGTTGTCCGGAATTATTGGGCGTA
AAGAGCTCGTAGGCGGCTTGTGCGCTCTGCTGTGAAAATCCGGGGCTTAACCTCCGGACCT
GCAGTGGATACGGGCAGGCTTGAGTTGCGTAGGGGAGACTGGAATTCCTGGTGTAGCGGT
GAAATGCGCAGATATCAGGAGGAACACCGGTGGCGAAGGCGGGTCTCTGGGCCGATACTG
ACGCTGAGGAGCGAAAGCGTGGGGAGCGAACAGGATTAGATACCCTGGTAGTCCACGCTG
TAAACGTTGGGCGCTAGGTGTGGGCGACTTCCACGTTGTCCGTGCCGTAGCTAACGCATT
AAGCGCCCCGCTGGGGAGTACGGCCGCAAGGCTAAAACTCAAAGGAATTGACGGGGGCC

>a1-1492R

TCCCTCCCCGCAAGCGGGGTTGGGCCATGGGCTTCGGGTGTTACCGACTTTCATGACGTG
ACGGGCGGTCTGTACAAAGGCCCGGAACGTATTACCGCAGCGTTGCTGATCTGCGATTA
CTAGCGACTCCGACTTCACGCAGTCGAGTTGCAGACTGCGATCCGAACTGAGACCGGCTT
TAAGGGATTTCGCTCCACCTCACGGTATCGCAGCCCTCTGTACCAGCCATTGTAGCATGTG
TGAAGCCCTGGACATAAGGGGCATGATGACTTGACGTCATCCCCACCTTCCTCCGAGTTG
ACCCCGGCAGTCTCCCGCGAGTCCCCGCCATTACGCGCTGGCAACACAGGACAAGGGTTG
CGCTCGTTGCGGGACTTAACCCAAACATCTCAGCACAGAGCTGACGACAGCCATGCACCA
CCTGCACACCGGCCACAAGGGGGCCGACATCTCTGCCGGTTTCCAGTGCATGTCAAGCCC
AGGTAAGGTTCTTCGCGTTGCATCGAATTAATCCACATGCTCCGCCGCTTGTGCGGGCCC
CCGTCAATTCCTTTGAGTTTTCGCTTGGCGCCGTACTCCCCAGGCGGGGCGCTTAATGC
GTTAGCTACGGCACGGACAACGTGGAACTCGCCCCACACCTAGCGCCCAACGTTTACAGCG
TGGACTACCAGGGTATCTAATCCTGTTTCGCTCCCCACGCTTTCGCTCCTCAGCGTCAGTA
TCGGCCCCAGAGACCCGCTTCGCCACCGGTGTTCCCTCCTGATATCTGCGCATTTACCGC
TACACCAGGAATTCAGTCTCCCCTACCGAACTCAAGCCTGCCCCGTATCCACTGCAGGTC
CGGAGTTAAGCCCCGGAT

>a1-1492R Inv compl

ATCCGGGGCTTAACCTCCGGACCTGCGAGTGGATACGGGCGAGGCTTGAGTTCCGGTAGGGGAGACTGGAATTCCTGGTG
TAGCGGTGAAATGCGCAGATATCAGGAGGAACACCGGTGGCGAAGGCGGGTCTCTGGGCCGATACTGACGCTGAGG
AGCGAAAGCGTGGGGAGCGAACAGGATTAGATACCCCTGGTAGTCCACGCTGTAAACGTTGGGCGCTAGGTGTGGGC
GACTTCCACGTTGTCCGTGCCGTAGCTAACGCATTAAGCGCCCCCGCTGGGGAGTACGGCCGCAAGGCTAAAACTC
AAAGGAATTGACGGGGGGCCGCAAGCGGGCGGAGCATTTGGATTAAATTCGATGCAACGCGGAAGAACCTTACCTGG
GCTTGACATGCACCTGGAACCCGGCAGAGATGTGGGCCCCCTTGTGTTCCGGTGTGCAGGTGGTTCATGGCTGTCTC
AGCTCGTGTGCTGAGATGTTGGGTTAAGTCCCGCAACGAGCGCAACCCCTTGTCTGTGTTGCCAGCGCGTAATGGC
GGGGACTCGCGGGAGACTGCCGGGGTCAACTCGGAGGAAGGTGGGGATGACGTCAGTCATCATGCCCTTATGTC
CAGGGCTTCACACATGCTACAAATGGCTGGTACAGAGGGCTGCGATACCGTGAGGTGGAGCGAATCCCTTAAAGCCG
GTCTCAGTTCGGATCGCAGTCTGCAACTCGACTGCGTGAAGTCCGGAGTCCCTAGTAATCGCAGATCAGCAACGCTG
CGGTGAATACGTTCCCGGGCTTGTACACACCGCCGTCACGTCATGAAAGTCGGTAACACCCGAAGCCCATGGCC
CAACCCCGCTTCCGGGGAGGGA

Seq A

GCTGAAGCACCTTCGGGTGTGGATGAGTGGCGAACGGGTGAGTAACACGTGGGTAACCTGCCCTGTACTTTGG
GATAAGCCCTGGAAACGGGGTCTAATACCGAATATGACTCGCTAAGGCATCTTGGTGGGTGGAAAGTTTCGGC
GGTACAGGATGGGCCCCGGGCTATCAGCTTCTTGGTGGGGTAGTGGCCTACCAAGCGGACGACGGGTAGCCG
GCCTGAGAGGGTGACCGGCCACACTGGGACTGAGACACGGCCCAGACTCCTACGGGAGGCAGCAGTGGGGAAT
ATTGCACAATGGGCGGAAGCCTGATGCAGCGACGCGCGGTGAGGGATGACGGCCTTCGGGTTGTAAACCTCTT
TCGCCAGGGACGAAGCGCAAGTGACGCTACCTGGAGAAGAAGCACCGGCTAACTACGTGCCAGCAGCCGCGGT
AATACGTTAGGGTGAAGCGTTGTCCGGAATTATTGGGCGTAAAGAGCTCGTAGGCGGCTTGTGCGCTCTGCTG
TGAAAATCCGGGGCTTAACCTCCGGACCTGCAGTGCATACGGGCAGGCTTGAGTTCCGGTAGGGGAGACTGGAAT
TCCTGGTGTAGCGGTGAAATGCGCAGATATCAGGAGGAACACCGGTGGCGAAGGCGGGTCTCTGGGCCGATAC
TGACGCTGAGGAGCGAAAGCGTGGGGAGCGAACAGGATTAGATACCCTGGTAGTCCACGCTGTAAACGTTGGG

CGCTAGGTGTGGGCGACTTCCACGTTGTCCGTGCCGTAGCTAACGCATTAAAGCGCCCCGCCTGGGGAGTACGG
CCGCAAGGCTAAAACTCAAAGGAATTGACGGGGGGCCCGCACAAAGCGGCGGAGCATGTGGATTAATTGGATGCAAC
GCGAAGAACCCTTACCTGGGCTTGACATGCACTGGAAACCGGCAGAGATGTGCGCCCCCTTGTGGCCGGTGTGCAGG
TGGTGCATGGGTGTGTCGTCAGCTCGTGTGCTGAGATGTTGGGTTAAGTCCCGCAACGAGCGCAACCCTTGTCTCTGTG
TTGCCAGCGCGTAATGGCGGGGACTCGCGGGAGACTGCCCGGGTCAACTCGGAGGAAAGGTGGGGATGACGTCAAGT
CATCATGCCCCCTTATGTCCAGGGCTTCACACATGCTACAATGGCTGGTACAGAGGGCTGCGATACCGTGAGGTGGA
GCGAATCCCTTAAAGCCGGTCTCAGTTCCGATCGCAGTCTGCAACTCGACTGCGTGAAGTCCGAGTCTGCTAGTAAT
CGCAGATCAGCAACGCTGCGGTGAATACGTTCCCGGGCTTGTACACACCGCCCTCACGTCATGAAAGTCGGTAA
CACCCGAAGCCCATGGCCCAACCCCGCTTGCGGGAGGGGA

>b2-27F

GCTGAAGCACCTTACGGGTGTGGATGAGTGGCGAACGGGTGAGTAACACGTGGGTAAACCT
GCCCTGTACTTTGGGATAAGCCCTGGAAACGGGGTCTAATACCGAATATGACTCGCTAAG
GCATCTTGGTGGGTGGAAAGTTTCGGCGGTACAGGATGGGCCCGCGGCCTATCAGCTTGT
TGGTGGGGTAGTGGCCTACCAAGGCGACGACGGGTAGCCGGCCTGAGAGGGTGACCGGCC
ACACTGGGACTGAGACACGGCCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGCAC
AATGGGCGGAAGCCTGATGCAGCGACGCCCGGTGAGGGATGACGGCCTTCGGGTTGTAAA
CCTCTTTTCGCCAGGGACGAAGCGCAAGTGACGGTACCTGGAGAAGAAGCACCGGCTAACT
ACGTGCCAGCAGCCGCGGTAATACGTAGGGTGCAAGCGTTGTCCGGAATTATTGGGCGTA
AAGAGCTCGTAGGCGGCTTGTCCGCTCTGCTGTGAAAAATCCGGGGCTTAACTCCGGACCT
GCAGTGGATACGGGCAGGCTTGAGTTCGGTAGGGGAGACTGGAATTCCTGGTGTAGCGGT
GAAATGCGCAGATATCAGGAGGAACACCGGTGGCGAAGGCGGGTCTCTGGGCCGATACTG
ACGCTGAGGAGCGAAAGCGTGGGGAGCGAACAGGATTAGATACCCTGGTAGTCCACGCTG
TAAACGTTGGGCGCTAGGTGTGGGCGACTTCCACGTTGTCCGTGCCGTAGCTAACGCATT
AAGCGCCCCGCCTTGGGGAGTACGGCCGCAAGGCTAAAACCTCAAAGGAATTGACGGGGGCC
CGCACAAAGC

>b2-1492R

CCCTCCCCGCAAGCGGGTTGGGCCATGGGCTTCGGGTGTTACCGACTTTCATGACGTGA
CGGGCGGTGTGTACAAGGCCCGGGAACGTATTACCGCAGCGTTGCTGATCTGCGATTAC
TAGCGACTCCGACTTCACGSCAGTCGAGTTGCAGACTGCGATCCGAACCTGAGACCGGCTTT
AAGGGATTTCGCTCCACCTCAGGSTATCGCAGCCCTCTGTACCAGCCATTGTAGCATGTGT
GAAGCCCTGGACATAAGGGGCATGATGACTTGACGTCATCCCCACCTTCCTCCGAGTTGA
CCCCGGCAGTCTCCCGCGAGTCCCGGCCATTACGCGCTGGCAACACAGGACAAGGGTTGC
GCTCGTTGCGGGACTTAACCCAACATCTCACGACACGAGCTGACGACAGCCATGCACCAC
CTGCACACCGGCCACAAGGGGGCCGACATCTCTGCCGGTTTCCAGTGCATGTCAAGCCCA
GGTAAGGTTCTTCGCGTTGCATCGAATTAATCCACATGCTCCGCCGCTTGTGCGGGCCCC
CGTCAATTCCCTTGAGTTTTCAGCTTGCGGCCGTACTCCCCAGGCGGGGCGCTTAATGCG
TTAGCTACGGCACGGACAACGTGGAAGTCGCCCACACCTAGCGCCCCAACGTTTACAGCGT
GGACTACAGGGTATCTAATCCCTGTTCCGCTCCCCACGCTTTCGCTCCTCAGCGTCAGTAT
CGGCCACGAGACCCGCCCTTCGCCACCGGTGTTCCCTCCTGATATCTGCGCATTTACCGCT
ACACCAGGAATTCCAGTCTCCCCCTACCGAACTCAAGCCTGCCCGTATCCACTGCAGGTC
CGGAGTTAAGCCCCGGAT

>b2-1492R

ATCCGGGGCTTAACTCCGGACCTGCAGTGGATACGGGCAGGCTTGAGTTGGGTAGGGGAGACTGGAATTCCTGGT
GTAGCGGTGAAATGCGCAGATATCAGGAGGAACACCGGTGGCGAAGGCGGGTCTCTGGGCGGATACCTGACGCTGAG
GAGCGAAAGCGTGGGGAGCGAACAGGATTAGATACCTTGGTAGTCCACGCTGTAAACGTTGGGCGCTAGGTGTGGG
CGACTTCCACGCTTGTCCGTGCGGTAGCTAAGCGCAATTAAGCGCCCCGCTGGGGAGTACGGCCGCAAGGCTTAAACT
CAAGGGAATTGACGGGGGGCCCGCACAAAGCGGGCGGAGCATGTGGATTAAATTCGATGCAACGCGGAAGAACCTTACCTG
GGCTTGACATGCACTGGAAACCGGCAGAGATGTCCGCCCGCTTGTGGCCGGTGTGCAGGTGGTGCATGGCTGTCT
CAGCTCGGTGTGCTGAGATGTTGGGTTAAGTCCCGCAACGAGCGCAACCTTGTGCTGTGTTGCCAGCGCGTAATGG
CGGGGACTCGCGGGAGACTGCCGGGCTCAACTCGGAGGAAGGTGGGGATGACGTCAAGTCATCATGCCCCCTTATGT
CCAGGGCTTCACACATGCTACAATGGCTGGTACAGAGGGCTGCGATACCGTGAGGTGGAGCGAATCCCTTAAAGCC
GGTCTCAGTTGGGATCGCAGTCTGCAACTCGACTGCGTGAAGTCGGAGTGGCTAGTAATCGCAGATCAGCAACGCT
GCGGTGAATAAGTTCCCGGGGCTTGTACACACCGCCCGTCAGCTCATGAAAGTCGGTAACACCCGAAGCCCATGGC
CCAACCCCGCTTGGCGGGAGGG

Seq B

GCTGAAGCACCTTCGGGTGTGGATGAGTGGCGAACGGGTGAGTAACACGTGGGTAAACCTGCCCTGTACTTTGG
GATAAGCCCTGGAAACGGGGTCTAATACCGAATATGACTCGCTAAGGCATCTTGGTGGGTGGAAAGTTTCGGC
GGTACAGGATGGGCCCGCGGCTATCAGCTTGTGGTGGGGTAGTGGCCTACCAAGGCGACGACGSGTAGCCG
GCCTGAGAGGGTGACCGGCCACACTGGGACTGAGACACGGCCCAGACTCCTACGGGAGGCAGCAGTGGGGAAT
ATTGCACAATGGGCGGAAGCCTGATGCAGCGACGCCCGGTGAGGGATGACGGCCTTCGGGTTGTAAACCTCTT
TCGCCAGGGACGAAGCGCAAGTGACGGTACCTGGAGAAGAAGCACCGGCTAACTACGTGCCAGCAGCCGCGGT

AATACGTAGGGTGCAAGCGTTGTCCGGAATTATTGGGCGTAAAGAGCTCGTAGGCGGCTTGTCGCGTCTGCTG
TGAAAAATCCGGGGCTTAAC TCCGGACCTGCACTGGATACGGGCAGGCTTGAGTTCCGTAGGGGAGACTGGAAT
TCCTGGTGTAGCGGTGAAATGCGCAGATATCAGGAGGAACACCGGTGGCGAAGGCGGGTCTCTGGGCGGATAC
TGACGCTGAGGAGCGAAAGCGTGGGGAGCGAACAGGATTAGATACCCCTGGTAGTCCACGCTGTAAACGTTGGG
CGCTAGGTGTGGGCGACTTCCACGTTGTCCGTGCCGTAGCTAACGCATTAAGCGCCCCGCCTGGGGAGTACGG
CCGCAAGGCTAAAACTCAAAGGAATTGACGGGGGGCCCGCACAAAGCGGCGGAGCATGTGGATTAATTCGATGCAA
CGCGAAGAACTTACCTGGGCTTGACATGCAC TGGAAACCGGCAGAGATGTGGGCCCCCTTGTTGGCCGGTGTGCAG
GTGGTGCATGGCTGTCTGTCAGCTCGTGTCTGTGAGATGTTGGGTAAAGTCCCGCAACGAGCGCAACCCCTTGTCCTGT
GTGGCCAGCGCGTAATGGCGGGGACTCGCGGGAGACTGCGGGGGTCAACTCGGAGGAAGGTGGGGATGACGTCAAG
TCATCATGCCCCCTTATGTCCAGGGCTTCACACATGCTACAATGGCTGGTACAGAGGGCTGGCGATACCGTGAGGTGG
AGCGAATCCCTTAAAGCCGGTCTCAGTTCCGATCGCAGTCTGCAACTCGACTGCGTGAAGTCCGGAGTCGCTAGTAA
TCGCAGATCAGCAACGCTGCGGTGAATACGTTCCCGGGCTTGTACACACCGCCCGTCACGTCATGAAAGTCGGTA
ACACCCGAAGCCCATGGCCCAACCCCGCTTGCGGGGAGGG

>c1-27F

NCTGAAGCACCTTCGGGTGTGGATGAGTGGCGAACGGGTGAGTAACACGTGGGTAACCTG
CCCTGTACTTTGGGATAAGCCCTGGAAACGGGGTCTAATACCGAATATGACTCGCTAAGG
CATCTTGGTGGGTGGAAAGTTTCGGCGGTACAGGATGGGCCCGCGGCCTATCAGCTTGTT
GGTGGGGTAGTGGCTTACCAAGGCGACGACGGGTAGCCGGCCTGAGAGGGTGACCGGCCA
CACTGGGACTGAGACACGGCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGCACA
ATGGGCGGAAGCCTGATGCAGCGACGCCGCTGAGGGATGACGGCCTTCGGGTGTAAAC
CTCTTTTCGCCAGGGACGAAGCGCAAGTGACGGTACCTGGAGAAGAAGCACCGGCTAACTA
CGTGCCAGCAGCCCGGTAATACGTAGGGTGCAAGCGTTGTCCGGAATTATTGGGCGTAA
AGAGCTCGTAGGCGGCTTGTGCGCTCTGCTGTGAAAATCCGGGGCTTAACGCCGACCTG
CAGTGGATACGGGCAGGCTTGAGTTCGGTAGGGGAGACTGGAATTCTGGGTGTAGCGGTG
AAATGCGCAGATATCAGGAGGAACACCGGTGCCGAAGGCGGCTCTCTGGGCCGATACTGA
CGCTGAGGAGCGAAAGCGTGGGGAGCGAACAGGATTAGATACCTGGTAGTCCACGCTGT
AAACGTTGGGCGCTAGGTGTGGGCGACTTCCACGTTGTCCGTGCCGTAGCTAACGCATTA
AGCGCCCCCGCTGGGGAGTACGGCCGCAAGGCTAAAACTCAAAGGAATTGACGGGGGCC

>c1-1492R

CCCTCCCCGCAAGCGGGGTGGGGCCATGGGCTTCGGGTGTTACCGACTTTCATGACGTGA
CGGGCGGTGTGTACAAGGCCCGGGAACGTATTACCGCAGCGTTGCTGATCTGCGATTAC
TAGCGACTCCGACTTCACGCAGTCGAGTTGCAGACTGCGATCCGAACCTGAGACCGGCTTT
AAGGGATTGCTCCACCTCAOGGTATCGCAGCCCTCTGTACCAGCCATTGTAGCATGTGT
GAAGCCCTGGACATAAGGGGCAATGACTTGACGTTCATCCCCACCTTCTCCGAGTTGA
CCCCGGCAGTCTCCCGCGAGTCCCCCGCATTACCGGCTGGCAACACAGGACAAGGGTTGC
GCTCGTTGCGGGGACTTAACCCAACATCTCACGACACGAGCTGACGACAGCCATGCACCAC
CTGCACACCGGCCACAAGGGGGCCGACATCTCTGCCGCTTTCAGTGCATGTCAAGCCCA
GGTAAGGTTCTTCGCGTTGCATCGAATTAATCCACATGCTCCGCCGCTTGTGCGGGCCCC
CGTCAATTCTTTGAGTTTGTAGCCTTTCGGCGCGTACTCCCCAGGCGGGGCGCTTAATGCG
TTAGCTACGGCACGGACAACGTGGAAGTCGCCACACCTAGCGCCCAACGTTTACAGCGT
GGACTACCAGGGTATCTAATCCTGTTGCTCCCCACGCTTTCGCTCCTCAGCGTCAGTAT
CGGCCACAGACCCGCTTCGCCACCGGTGTTCCTCCTGATATCTGCGCATTTACCGCT
ACACCAGGAATTCAGTCTCCCCCTACCGAACTCAAGCCTGCCCCGTATCCACTGCAGGTCC
GGAGTTAAGCCCCGGAT

>c1-1492R

ATCCGGGGCTTAACCTCCGGACCTGCAGTGGATACGGGCAGGCTTGASTTCGGTAGGGGAGACTGGAATTCCTGGTGTAGC
GGTGAAATGCGCAGATATCAGGAGGAACACCGGTGGCGAAGGCGGGTCTCTGGGCCGATACTGACGCTGAGGAGCGAAAG
CGTGGGGAGCGAACAGGATTAGATACCTTGGTAGTCCACGCTGTAAACGTTGGGCGCTAGGTTGTGGGCGACTTCCACGTT
GTCCGTGCCGTAGCTAACGCATTAAGCGCCCCCGCTGGGGAGTACGGCCGCAAGGCTAAAACTCAAAGGAATTGACGGGG
GCCCGCACAAAGCGCGGAGCATGTGATTAATTCGATGCAACGCGAAGAACCCTTACCTGGGCTTGACATGCAGTGGAAAC
CGGCAGAGATGTGCGCCCCCTTGTGGCCGGTGTGCAGGTGGTGCATGGCTGTCTGCTCAGCTCGTGTCTGAGATGTTGGGT
TAAGTCCCAGCAACGAGCGCAACCTTGTCTGTGTTGCCAGCGCTAATGGCGGGGACTCGCGGGAGACTGCCGGGGTCA
ACTCGGAGGAAGGTGGGGATGACGTCAAGTCAATCATGCCCTTATGTCCAGGGCTTCACACATGCTACAATGGCTGGTAC
AGAGGGCTGCGATACCGTGAGGTGGAGCGAATCCCTTAAAGCCGGTCTCAGTTTCGGATCGCAGTCTGCAACTCGACTGCG
TGAAGTGCGAGTTCGTAGTAATCGCAGATCAGCAACGCTGCGGTGAATACGTTCCCGGGCCCTGTACACACCGCCCGTCA
CGTCATGAAAGTCGGTAACACCCGAAGCCCCATGGCCCCAACCCCGCTTTCGGGGAGGG


Seq C

NCTGAAGCACCTTCGGGTGTGGATGAGTGGCGAACGGGTGAGTAACACGTGGGTAACCTGCCCTGTACTTTGG
GATAAGCCCTGGAAACGGGGTCTAATACCGAATATGACTCGCTAAGGCATCTTGGTGGGTGGAAAGTTTCGGC
GGTACAGGATGGGCCCGCGGCTATCAGCTTGTTCGGTGGGGTAGTGGCTTACCAAGGCGACGACGGGTAGCCG
GCCTGAGAGGGTGAACCGGCCACATGGGACTGAGACACGGCCAGACTCCTACGGGAGGCAGCAGTGGGGAAT
ATTGCACAATGGGCGGAAGCCTGATGCAGCGACGCCGCTGAGGGATGACGGCCTTCGGGTGTAAACCTCTT
TCGCCAGGGACGAAGCGCAAGTGACGGTACCTGGAGAAGAAGCACCGGCTAACTACGTGCCAGCAGCCGCGGT
AATACGTAGGGTGCAAGCGTTGTCCGGAATTATTGGGCGTAAAGAGCTCGTAGGCGGCTTGTGCGCTCTGCTG
TGAAAATCCGGGGCTTAACCTCCGGACCTGCAGTGGATACGGGCAGGCTTGAGTTCGGTAGGGGAGACTGGAAT

TCCTGGTGTAGCGGTGAAATGCCGAGATATCAGGAGGAACACCGGTGGCGAAGGCGGGTCTCTGGGCGGATAC
TGACGCTGAGGAGCGAAAGCGTGGGGAGCGAACAGGATTAGATACCCCTGGTAGTCCACGCTGTAAACGTTGGG
CGCTAGGTGTGGGCGACTTCCACGTTGTCCGTGCCGTAGCTAACGCATTAAAGCGCCCCGCCCTGGGGAGTACGG
CCGCAAGGCTAAAACCTCAAAGGAATTGACGGGGGCCCCGCACAAGCGCGGAGCATGTGGATTAATTCGATGCAACG
CGAAGAACCTTACCTGGGCTTGACATGCACTGGAAACCGGCAGAGATGTCGGCCCCCTTGTGGCCGCTGTGCAGGTGGTG
CATGGCTGTGTCAGCTCGTGTCTGAGATGTTGGGTAAAGTCCCACAACGAGCGCAACCCCTGTCTGTGTTGCCAGCG
CGTAATGGCGGGGACTCGCGGGGAGACTGCCGGGGTCAACTCGGAGGAAGGTGGGGATGACGTCAAGTCATCATGCCCTT
ATGTCCAGGGCTTCACACATGCTACAATGGCTGGTACAGAGGGCTGCGATACCGTGAGGTGGAGCGAATCCCTTAAAGCC
GGTCTCAGTTCGGATCGCAGTCTGCAACTCGACTGCGTGAAGTCGGAGTCGCTAGTAATCGCAGATCAGCAACGCTGCGG
TGAATACGTTCCCGGGCCTTGTACACACCGCCCGTCACGTCATGAAAGTCGGTAACACCCGAAGCCCATGGCCCAACCCC
GCTTGCGGGGAGGG

0.02

Exhibit 5



EXTERNAL MEMORANDUM

Re : Report of production trials according to US 6,133,003 to Rabenhorst et al.

Table of contents

1. Goal.....	2
2. Material and Methods.....	2
2.1 Cultures work out.....	2
2.1.1 Inoculation chain.....	2
2.1.2 Cultures in fermenters (max. working volume 15L).....	3
2.2 Cultures follow up.....	5
2.2.1 Culture parameters.....	5
2.2.2 Bacteriological follow up.....	5
2.2.3 HPLC Analyses	6
3. Results.....	6
3.1 Culture follow up.....	6
3.1.1 DSM 9992.....	6
3.1.2 Zyl 926 (Asp L-ii 2).....	8
3.1.3 Cultures comparison.....	9
3.2 Analytical Results	9
4. Conclusion.....	12

1. Goal

The goal is to compare two strains :

- *Amycolatopsis* sp. DSM 9992
- *Amycolatopsis* sp. Zyl 926 (referred to also as Asp L-ii 2)

when cultivated in identical conditions, namely identical inoculating chain, fermentation process and equipment.

These two strains were evaluated for the production of vanillin by reducing to practice the process taught in US Patent 6,133,003 to Rabenhorst et al, sometimes referred to as the Symrise process in the following. At the end of the present memo, some results are given while using strain Zyl 926 (Asp Lii-2) according to the process taught in US Application N° 10/550,945 to Heald et al, referred to as the Safisis process.

2. Material and Methods

2.1 Cultures work out

The process applied in this study was according to the teaching of US Patent 6,133,003 to Rabenhorst et al "Process for the preparation of vanillin and microorganism suitable therefor".

2.1.1. Inoculating chain

- According to working Example 1 of US 6,133,003

Preparation of the preliminary culture

*A 500 ml conical flask with a side baffle was filled with 100 ml of medium, comprising 1 g of malt extract, 0,4 g of glucose and 0,4 g of yeast extract and made up to 100 ml with water and then sterilized for 20 min at 121°C. After cooling, the flask was inoculated with 200 µl of a frozen glycerol culture of *Amycolatopsis* sp. DSM 9992 or Zyl 926 (Asp L-ii 2). The cultures were incubated on a rotary shaking machine at 45 °C and 100 rpm. After 24 h, these cultures were used to inoculate the production medium.*

- Preliminary culture medium preparation

Product	Concentration
Glucose	4 g/Kg
Malt extract	10 g/Kg
Yeast extract	4 g/Kg
H ₂ O	q.s.p
pH not adjusted	
Sterilization 20' at 121°C	

- Preculture inoculation

*Inoculation of 100 mL with 0,2 mL from the content of a cryotube (frozen glycerol culture)

*Incubation (shaking machine) : 45°C - 100 rpm - 24h

2.1.2 Cultures in fermenters (maximum working volume = 15L)

- Stage production according to working Example 3 of US 6,133,003

Vanillin Production in a 10 L fermenter.

5 l of culture medium (4 g/l of glucose, 10 g/l of malt extract and 6 g/l of yeast extract) were sterilized in a fermenter and, after cooling, were inoculated with 100 ml of a seed culture of DSM 9992 or Zyl 926 (Asp L-ii 2) according to Example 1.

According to working Example 3, the culture conditions were : 37°C, 500 rpm, 5 l of air/min, 12,5 h after inoculation, 1.634 kg of an approximately 3.7% strength ferulic acid solution (60.2 g of ferulic acid) were added. After 17.5 h, a further 4.397 kg of an approximately 3.7% strength ferulic acid solution (164.72 g of ferulic acid) were pumped in over a period of 10 h.

While trying to reproduce these very same parameters, I had to adapt some of the above data to make it work and arrive at the following parameters :

- Culture parameters

	GROWTH	BIOCONVERSION	BIOCONVERSION
Inoculation	100 mL	-	-
Temperature	37°C	37°C	37°C
Medium quantity	5 Kg	1.6 Kg	4.4 Kg
Culture condition	Batch	Batch (isolated addition of ferulic acid)	Fed-batch during 10 h (7.2 mL/min)
Duration	12.5h	5h	14.5h
pO ₂ Regulation	50% (not mentioned in the example 3 of the patent)		
Aeration	5 L/min	7 L/min	7 L/min
Stirring	400 à 750 rpm		
pH regulation	8 ± 0.5 (not mentioned in the example 3 of the patent but in the spec.)		
Surpression	No		
End of culture	-	-	32 h

- Fermenters preparation

Product	Concentration
Glucose	4 g/Kg
Malt extract	10 g/Kg
Yeast extract	6 g/Kg
Propylene glycol P2000	0.35 g/Kg
H ₂ O	q.s.p
5 Kg of medium in the fermenter <i>in situ</i> sterilization	

- Ferulic acid solution

Product	Concentration
Ferulic acid	3.75%
pH adjustment : 7.4 with a sodium hydroxyde solution*	

*preliminary tests to dissolve ferulic acid in water were carried out in vain. I had to add sodium hydroxide otherwise I could not get any homogeneous solution.

Detailed sequence of addition of the ferulic acid solution :

- At T12,5 h of culture, 1,6 kg of the ferulic solution were added in the fermenter

- For the semi-continuous step which starts at 17,5 h of culture, 1,3 kg of solution were prepared to feed in the fermenter. It was necessary to reload the solution every 2,5 h because a degradation of ferulic acid is observed when the solution is prepared more than 3 h in advance. The flow rate is 7,2 ml/min during 10 h, allowing to add 4,4 kg of ferulic acid solution.

- After 32 h, the fermentation was terminated.

2.2 Culture follow up

2.2.1 Culture parameters

For each culture, parameters were monitored as follows :

- pO₂ (dissolved oxygen in the medium) which allows to visualize the bacterial growth by the more or less important oxygen demand. This parameter is controlled by regulating aeration and/or agitation of the medium to a certain level in order to secure optimal growth conditions
- pH is regulated to favor bacterial growth and the production of metabolites of interest
- Solutions used for pH regulation [sulfuric acid (H₂SO₄) and/or sodium hydroxyde (NaOH)]

2.2.2 Bacteriological follow up

A bacterial follow up was realized in order to be ascertain the absence of any contamination.

2.2.3 HPLC Analyses

For each culture, several samples were withdrawn at mentioned times and frozen until analyzed :

T0
T4
T8
T12,5 after spot addition of ferulic acid
T17,5 after the starting of the semi-continuous culture
T20 before reloading of the ferulic acid solution
T22,5 before reloading of the ferulic acid solution
T25 before reloading of the ferulic acid solution
T27,5 before termination of the semi-continuous culture
T32

Production of vanillin, vanillic acid, vanillic alcohol, guaiacol and of the ferulic acid consumption were assayed by HPLC (high pressure liquid chromatography) and UV detection, consumption of glucose was tested through HPLC and refractometry.

Two dilutions were made for each sample in order to be sure to be in the detection range of the metabolites.

3. Results

3.1 Culture follow up

3.1.1 DSM 9992

As illustrated by Figure 1, growth phase went fine, as shown by the pO₂ which decreased from 100 % to 57 % between T0 and T12,5 h. This pO₂ value had slightly risen during the spot addition of ferulic acid (to 61 %) then decreased again regularly to reach 36 % at the termination of the culture. The 50 % instruction value for pO₂ could not be maintained, though agitation was set at a high limit but pO₂ did not drop under 30 %, thus oxygen supply has been sufficient throughout the culture duration.

Regarding pH regulation, the addition of acid and/or base solution was not necessary, let alone at the starting of the culture to restore the pH at $8,0 \pm 0,5$.

At T24 h, I observed a high foam production necessitating the addition of antifoam propylene glycol P2000.

At the termination of the culture, the culture medium was of a yellow/orange color, greasy –probably due to the presence of antifoam) and I noticed very high levels of clove odor/spice, showing *a priori* that bioconversion practically did not occur.

SYMRISE process : parameter acquisition (February 2010)
DSM 9992 strain





Figure 1 : Parameter acquisition for DSM 9992 strain cultivated according to Symrise process (US 6,133,033) (Cumul = Total)

3.1.2 Zyl 926 (Asp L-ii 2)

As shown in Figure 2, this experiment took place with noticeable differences compared to the previous one with strain DSM 9992 :

- pO₂ decreased regularly during growth phase then went over 80 % after the spot addition of ferulic acid
- pH regulation by base solution addition only after addition of ferulic acid during the bioconversion (total amount added : 120g of NaOH 10%).

SYMRISE process : parameter acquisition (February 2010)
Asp L-ii 2 strain





Figure 2 : Parameter acquisition for the Zyl 926 (Asp L-ii 2) strain cultivated according to the Symrise process (US 6,133,033)

3.1.3 Cultures comparison

Table 1 illustrates the principal data :

Table 1 : Growth comparison of DSM 9992 and Zyl 926 (Asp L-ii 2) strains

	pO ₂	pH	pH regulation with sodium hydroxide 10%	pH regulation with sulfuric acid 10%	Medium composition at the end of the culture
DSM 9992	Growth phase (between 0 - 12 h): decrease from 100% to 57% Bioconversion phase (between 12 - 32 h) : increase to 61% after ferulic acid addition, then decrease until the end (36% at 32h)	Low initial pH (5.4) → addition of 15g of sodium hydroxide 10% to adjust pH at 8.0 ± 0.5	Bioconversion phase (between 12 - 32 h) : NO	NO	Oily medium (due to antifoam), yellow/orange, strong smell like a clove
Zyl 926 (Asp L-ii 2)	Growth phase (between 0 - 12 h): continuous decrease from 100% to 69% Bioconversion phase (between 12 - 32 h) : increase to 85% after ferulic acid addition, then static until the end	No need to adjust initial pH	Continuous addition of sodium hydroxide 10% during bioconversion phase (quantity : 120g)	NO	Orange medium, smell like butter

3.2 Analytical results

Table 2 allows to compare the results after the termination of the culture for the two strains :

Table 2 : End product concentrations for Zyl 926 (Asp L-ii 2) and DSM 9992 strains when cultivated according to Example 3 of US 6,133,003

Strain	Vanillic alcohol (g/Kg)	Vanillic acid (g/Kg)	Vanillin (g/Kg)	Guaiacol (g/Kg)
Asp L-ii 2	0	0	0	0
DSM 9992	<0.1	0.3	<0.1	0

Chromatograms below clearly indicate that there was an accumulation of ferulic acid over time, whatever strain was used.

No metabolite of interest was detected for strain Zyl 926 (Asp L-ii 2) as illustrated in Figure 3 below, and as shown in Figure 4 below, almost none in the case of strain DSM 9992 : under 0,1 g/kg (0,1 g/l) regarding vanillin, which value should be interpreted as follows : practically no vanillin was produced contrary to the statement in Example 3 of US 6,133,003.

Moreover, at T17,5 h-T20 h-T22,5 h-T25 h and T27,5 h, analysis of intermediary samples was conducted and showed no evidence of the presence of metabolites which could have been produced and then eventually degraded during the cultivation period.




Figure 3 : Chromatogram/Symrise process (US 6,133033) with Zyl 926 strain (Asp L-ii 2) -- sample corresponding to the end of the culture (T32 h)

[French/English translation: alcool vanillique/vanillic alcohol -- acide vanillique/vanillic acid -- vanilline/vanillin -- guaiacol/guaiacol -- acide férulique/ferulic acid]




Figure 4 : Chromatogram/Symrise process (US 6,133,033) with DSM 9992 strain – sample corresponding to the end of the culture (T32 h)

[French/English translation: alcoo1 vanillique/vanillic alcohol – acide vanillique/vanillic acid – vanilline/vanillin – guaiacol/guaiacol – acide férulique/ferulic acid]

Table 3 illustrates the accumulation profile of ferulic acid during cultures.

Table 3 : Accumulation profile of ferulic acid in the culture medium

Sample	Ferulic acid Theoretical quantity added (g)	Ferulic acid Theoretical quantity cumulated* (g)	Ferulic acid Theoretical concentration* (g/Kg)	Ferulic acid HPLC results (g/Kg)
T12.5 after addition of ferulic acid solution	60.2 g	-	9.1 g/Kg	8.9 (± 0.3)**
T20 before fed-batch reloading	48.75 g	108.9 g	13.8 g/kg	15.0 (± 2.4)**
T22.5 before fed-batch reloading	48.75 g	157.7 g	17.1 g/Kg	17.3 (± 1.8)**
T25 before fed-batch reloading	48.75 g	206.5 g	19.7 g/Kg	18.5 (± 1.6)**
T27.5 at the end of fed-batch culture	48.75 g	255.3 g	21.6 g/Kg	21.1 (± 0.8)**
T32				19.8 (± 1.1)**

*Two hypothesis :

- no addition of sodium hydroxide to regulate pH
- no consumption and/or loss of ferulic acid between fed-batch reloading

** Result average (g/Kg)

As illustrated by theoretical calculations reported in Table 3 above, bioconversion phase was conducted as taught in US patent 6,133,003 but ferulic acid accumulated and its consumption and degradation after introduction in the culture medium are negligible.

4. Conclusion

Though conducted with identical culture conditions, the above analytical results allow to conclude that strain DSM 9992, claimed in US 6,133,003 and Zyl 926 (Asp L-ii 2), claimed in US Application N° 10/550,945, have shown a different behavior when grown in fermenters. Besides, in both cases growth phase went well.

Moreover, HPLC analysis were performed in order to quantify metabolites of interest, namely vanillin, guaiacol, vanillic alcohol and vanillic acid expected to be produced in such conditions.

These trials results allow us to conclude that no ferulic acid reacted and the latter accumulated in the culture medium during the course of the culture.

These were unexpected results since the parameters applied in the present study were those taught in working Examples of the document US 6,133,003 to Rabenhorst et al., "Process for the preparation of vanillin and microorganisms suitable therefor".

Moreover we, present engineers who signed the appended affidavits, made all reasonable attempts to use the complete teaching of this specification and not only the Example wording. See for example to name but a few "adjustments", the preparation of ferulic acid and also pH and pO₂ regulation and antifoam necessary usage.

As a conclusion, and based on the forgoing facts, this prior art, namely US 6,133,003 is of a non operative kind, the reason being that a person skill in the art by repeating/reproducing its teaching not only does not get the expected results but rather gets essentially no production of the desired product, namely vanillin. To arrive at this production, one should "invent" another process.

Accordingly, US 6,133,003 is not at all pertinent, and by far, the production of vanillin as taught and claimed by the inventors of US Application N° 10/550,945 is not obviously derivable therefrom.

By way of comparison below is shown in Figure 5, a chromatogram obtained by working out the process taught in US Application N° 10/550,945 to Heald et al, namely the "Safisis process", using Zyl 926 (Asp L-ii 2) : ferulic acid was spent and the vanillin peak is predominant (14 g/l).




Figure 5 : Chromatogram/Safisis process (US App. N° 10/550,945) with Zyl 926 strain (Asp L-ii 2) - sample withdrawn at the end of the culture

[French/English translation: alcool vanillique/vanillic alcohol – acide vanillique/vanillic acid – vanilline/vanillin – guaiacol/guaiacol – acide férulique/ferulic acid]